

**PATENT**

Attorney Docket No.: 119927-1030

**METHODS AND APPARATUS FOR IDENTIFYING ALLOSTERICALLY  
REGULATED RIBOZYMES**

5

**FIELD OF THE INVENTION**

The present invention relates generally to the field of  
ribozymes and in particular to aptazymes or allosteric,  
regulatable ribozymes that modulate their kinetic parameters  
in response to the presence of an effector molecule.

10  
11  
12  
13  
14  
15  
16  
17  
18  
19  
20  
21  
22  
23  
24  
25  
26  
27  
28  
29  
30  
31  
32  
33  
34  
35  
36  
37  
38  
39  
40  
41  
42  
43  
44  
45  
46  
47  
48  
49  
50  
51  
52  
53  
54  
55  
56  
57  
58  
59  
60  
61  
62  
63  
64  
65  
66  
67  
68  
69  
70  
71  
72  
73  
74  
75  
76  
77  
78  
79  
80  
81  
82  
83  
84  
85  
86  
87  
88  
89  
90  
91  
92  
93  
94  
95  
96  
97  
98  
99  
100  
101  
102  
103  
104  
105  
106  
107  
108  
109  
110  
111  
112  
113  
114  
115  
116  
117  
118  
119  
120  
121  
122  
123  
124  
125  
126  
127  
128  
129  
130  
131  
132  
133  
134  
135  
136  
137  
138  
139  
140  
141  
142  
143  
144  
145  
146  
147  
148  
149  
150  
151  
152  
153  
154  
155  
156  
157  
158  
159  
160  
161  
162  
163  
164  
165  
166  
167  
168  
169  
170  
171  
172  
173  
174  
175  
176  
177  
178  
179  
180  
181  
182  
183  
184  
185  
186  
187  
188  
189  
190  
191  
192  
193  
194  
195  
196  
197  
198  
199  
200  
201  
202  
203  
204  
205  
206  
207  
208  
209  
210  
211  
212  
213  
214  
215  
216  
217  
218  
219  
220  
221  
222  
223  
224  
225  
226  
227  
228  
229  
230  
231  
232  
233  
234  
235  
236  
237  
238  
239  
240  
241  
242  
243  
244  
245  
246  
247  
248  
249  
250  
251  
252  
253  
254  
255  
256  
257  
258  
259  
260  
261  
262  
263  
264  
265  
266  
267  
268  
269  
270  
271  
272  
273  
274  
275  
276  
277  
278  
279  
280  
281  
282  
283  
284  
285  
286  
287  
288  
289  
290  
291  
292  
293  
294  
295  
296  
297  
298  
299  
300  
301  
302  
303  
304  
305  
306  
307  
308  
309  
310  
311  
312  
313  
314  
315  
316  
317  
318  
319  
320  
321  
322  
323  
324  
325  
326  
327  
328  
329  
330  
331  
332  
333  
334  
335  
336  
337  
338  
339  
340  
341  
342  
343  
344  
345  
346  
347  
348  
349  
350  
351  
352  
353  
354  
355  
356  
357  
358  
359  
360  
361  
362  
363  
364  
365  
366  
367  
368  
369  
370  
371  
372  
373  
374  
375  
376  
377  
378  
379  
380  
381  
382  
383  
384  
385  
386  
387  
388  
389  
390  
391  
392  
393  
394  
395  
396  
397  
398  
399  
400  
401  
402  
403  
404  
405  
406  
407  
408  
409  
410  
411  
412  
413  
414  
415  
416  
417  
418  
419  
420  
421  
422  
423  
424  
425  
426  
427  
428  
429  
430  
431  
432  
433  
434  
435  
436  
437  
438  
439  
440  
441  
442  
443  
444  
445  
446  
447  
448  
449  
450  
451  
452  
453  
454  
455  
456  
457  
458  
459  
460  
461  
462  
463  
464  
465  
466  
467  
468  
469  
470  
471  
472  
473  
474  
475  
476  
477  
478  
479  
480  
481  
482  
483  
484  
485  
486  
487  
488  
489  
490  
491  
492  
493  
494  
495  
496  
497  
498  
499  
500  
501  
502  
503  
504  
505  
506  
507  
508  
509  
510  
511  
512  
513  
514  
515  
516  
517  
518  
519  
520  
521  
522  
523  
524  
525  
526  
527  
528  
529  
530  
531  
532  
533  
534  
535  
536  
537  
538  
539  
540  
541  
542  
543  
544  
545  
546  
547  
548  
549  
550  
551  
552  
553  
554  
555  
556  
557  
558  
559  
560  
561  
562  
563  
564  
565  
566  
567  
568  
569  
570  
571  
572  
573  
574  
575  
576  
577  
578  
579  
580  
581  
582  
583  
584  
585  
586  
587  
588  
589  
590  
591  
592  
593  
594  
595  
596  
597  
598  
599  
600  
601  
602  
603  
604  
605  
606  
607  
608  
609  
610  
611  
612  
613  
614  
615  
616  
617  
618  
619  
620  
621  
622  
623  
624  
625  
626  
627  
628  
629  
630  
631  
632  
633  
634  
635  
636  
637  
638  
639  
640  
641  
642  
643  
644  
645  
646  
647  
648  
649  
650  
651  
652  
653  
654  
655  
656  
657  
658  
659  
660  
661  
662  
663  
664  
665  
666  
667  
668  
669  
670  
671  
672  
673  
674  
675  
676  
677  
678  
679  
680  
681  
682  
683  
684  
685  
686  
687  
688  
689  
690  
691  
692  
693  
694  
695  
696  
697  
698  
699  
700  
701  
702  
703  
704  
705  
706  
707  
708  
709  
710  
711  
712  
713  
714  
715  
716  
717  
718  
719  
720  
721  
722  
723  
724  
725  
726  
727  
728  
729  
730  
731  
732  
733  
734  
735  
736  
737  
738  
739  
740  
741  
742  
743  
744  
745  
746  
747  
748  
749  
750  
751  
752  
753  
754  
755  
756  
757  
758  
759  
760  
761  
762  
763  
764  
765  
766  
767  
768  
769  
770  
771  
772  
773  
774  
775  
776  
777  
778  
779  
780  
781  
782  
783  
784  
785  
786  
787  
788  
789  
790  
791  
792  
793  
794  
795  
796  
797  
798  
799  
800  
801  
802  
803  
804  
805  
806  
807  
808  
809  
810  
811  
812  
813  
814  
815  
816  
817  
818  
819  
820  
821  
822  
823  
824  
825  
826  
827  
828  
829  
830  
831  
832  
833  
834  
835  
836  
837  
838  
839  
840  
841  
842  
843  
844  
845  
846  
847  
848  
849  
850  
851  
852  
853  
854  
855  
856  
857  
858  
859  
860  
861  
862  
863  
864  
865  
866  
867  
868  
869  
870  
871  
872  
873  
874  
875  
876  
877  
878  
879  
880  
881  
882  
883  
884  
885  
886  
887  
888  
889  
890  
891  
892  
893  
894  
895  
896  
897  
898  
899  
900  
901  
902  
903  
904  
905  
906  
907  
908  
909  
910  
911  
912  
913  
914  
915  
916  
917  
918  
919  
920  
921  
922  
923  
924  
925  
926  
927  
928  
929  
930  
931  
932  
933  
934  
935  
936  
937  
938  
939  
940  
941  
942  
943  
944  
945  
946  
947  
948  
949  
950  
951  
952  
953  
954  
955  
956  
957  
958  
959  
960  
961  
962  
963  
964  
965  
966  
967  
968  
969  
970  
971  
972  
973  
974  
975  
976  
977  
978  
979  
980  
981  
982  
983  
984  
985  
986  
987  
988  
989  
990  
991  
992  
993  
994  
995  
996  
997  
998  
999  
1000  
1001  
1002  
1003  
1004  
1005  
1006  
1007  
1008  
1009  
1010  
1011  
1012  
1013  
1014  
1015  
1016  
1017  
1018  
1019  
1020  
1021  
1022  
1023  
1024  
1025  
1026  
1027  
1028  
1029  
1030  
1031  
1032  
1033  
1034  
1035  
1036  
1037  
1038  
1039  
1040  
1041  
1042  
1043  
1044  
1045  
1046  
1047  
1048  
1049  
1050  
1051  
1052  
1053  
1054  
1055  
1056  
1057  
1058  
1059  
1060  
1061  
1062  
1063  
1064  
1065  
1066  
1067  
1068  
1069  
1070  
1071  
1072  
1073  
1074  
1075  
1076  
1077  
1078  
1079  
1080  
1081  
1082  
1083  
1084  
1085  
1086  
1087  
1088  
1089  
1090  
1091  
1092  
1093  
1094  
1095  
1096  
1097  
1098  
1099  
1100  
1101  
1102  
1103  
1104  
1105  
1106  
1107  
1108  
1109  
1110  
1111  
1112  
1113  
1114  
1115  
1116  
1117  
1118  
1119  
1120  
1121  
1122  
1123  
1124  
1125  
1126  
1127  
1128  
1129  
1130  
1131  
1132  
1133  
1134  
1135  
1136  
1137  
1138  
1139  
1140  
1141  
1142  
1143  
1144  
1145  
1146  
1147  
1148  
1149  
1150  
1151  
1152  
1153  
1154  
1155  
1156  
1157  
1158  
1159  
1160  
1161  
1162  
1163  
1164  
1165  
1166  
1167  
1168  
1169  
1170  
1171  
1172  
1173  
1174  
1175  
1176  
1177  
1178  
1179  
1180  
1181  
1182  
1183  
1184  
1185  
1186  
1187  
1188  
1189  
1190  
1191  
1192  
1193  
1194  
1195  
1196  
1197  
1198  
1199  
1200  
1201  
1202  
1203  
1204  
1205  
1206  
1207  
1208  
1209  
1210  
1211  
1212  
1213  
1214  
1215  
1216  
1217  
1218  
1219  
1220  
1221  
1222  
1223  
1224  
1225  
1226  
1227  
1228  
1229  
1230  
1231  
1232  
1233  
1234  
1235  
1236  
1237  
1238  
1239  
1240  
1241  
1242  
1243  
1244  
1245  
1246  
1247  
1248  
1249  
1250  
1251  
1252  
1253  
1254  
1255  
1256  
1257  
1258  
1259  
1260  
1261  
1262  
1263  
1264  
1265  
1266  
1267  
1268  
1269  
1270  
1271  
1272  
1273  
1274  
1275  
1276  
1277  
1278  
1279  
1280  
1281  
1282  
1283  
1284  
1285  
1286  
1287  
1288  
1289  
1290  
1291  
1292  
1293  
1294  
1295  
1296  
1297  
1298  
1299  
1300  
1301  
1302  
1303  
1304  
1305  
1306  
1307  
1308  
1309  
1310  
1311  
1312  
1313  
1314  
1315  
1316  
1317  
1318  
1319  
1320  
1321  
1322  
1323  
1324  
1325  
1326  
1327  
1328  
1329  
1330  
1331  
1332  
1333  
1334  
1335  
1336  
1337  
1338  
1339  
1340  
1341  
1342  
1343  
1344  
1345  
1346  
1347  
1348  
1349  
1350  
1351  
1352  
1353  
1354  
1355  
1356  
1357  
1358  
1359  
1360  
1361  
1362  
1363  
1364  
1365  
1366  
1367  
1368  
1369  
1370  
1371  
1372  
1373  
1374  
1375  
1376  
1377  
1378  
1379  
1380  
1381  
1382  
1383  
1384  
1385  
1386  
1387  
1388  
1389  
1390  
1391  
1392  
1393  
1394  
1395  
1396  
1397  
1398  
1399  
1400  
1401  
1402  
1403  
1404  
1405  
1406  
1407  
1408  
1409  
1410  
1411  
1412  
1413  
1414  
1415  
1416  
1417  
1418  
1419  
1420  
1421  
1422  
1423  
1424  
1425  
1426  
1427  
1428  
1429  
1430  
1431  
1432  
1433  
1434  
1435  
1436  
1437  
1438  
1439  
1440  
1441  
1442  
1443  
1444  
1445  
1446  
1447  
1448  
1449  
1450  
1451  
1452  
1453  
1454  
1455  
1456  
1457  
1458  
1459  
1460  
1461  
1462  
1463  
1464  
1465  
1466  
1467  
1468  
1469  
1470  
1471  
1472  
1473  
1474  
1475  
1476  
1477  
1478  
1479  
1480  
1481  
1482  
1483  
1484  
1485  
1486  
1487  
1488  
1489  
1490  
1491  
1492  
1493  
1494  
1495  
1496  
1497  
1498  
1499  
1500  
1501  
1502  
1503  
1504  
1505  
1506  
1507  
1508  
1509  
1510  
1511  
1512  
1513  
1514  
1515  
1516  
1517  
1518  
1519  
1520  
1521  
1522  
1523  
1524  
1525  
1526  
1527  
1528  
1529  
1530  
1531  
1532  
1533  
1534  
1535  
1536  
1537  
1538  
1539  
1540  
1541  
1542  
1543  
1544  
1545  
1546  
1547  
1548  
1549  
1550  
1551  
1552  
1553  
1554  
1555  
1556  
1557  
1558  
1559  
1560  
1561  
1562  
1563  
1564  
1565  
1566  
1567  
1568  
1569  
1570  
1571  
1572  
1573  
1574  
1575  
1576  
1577  
1578  
1579  
1580  
1581  
1582  
1583  
1584  
1585  
1586  
1587  
1588  
1589  
1590  
1591  
1592  
1593  
1594  
1595  
1596  
1597  
1598  
1599  
1600  
1601  
1602  
1603  
1604  
1605  
1606  
1607  
1608  
1609  
1610  
1611  
1612  
1613  
1614  
1615  
1616  
1617  
1618  
1619  
1620  
1621  
1622  
1623  
1624  
1625  
1626  
1627  
1628  
1629  
1630  
1631  
1632  
1633  
1634  
1635  
1636  
1637  
1638  
1639  
1640  
1641  
1642  
1643  
1644  
1645  
1646  
1647  
1648  
1649  
1650  
1651  
1652  
1653  
1654  
1655  
1656  
1657  
1658  
1659  
1660  
1661  
1662  
1663  
1664  
1665  
1666  
1667  
1668  
1669  
1670  
1671  
1672  
1673  
1674  
1675  
1676  
1677  
1678  
1679  
1680  
1681  
1682  
1683  
1684  
1685  
1686  
1687  
1688  
1689  
1690  
1691  
1692  
1693  
1694  
1695  
1696  
1697  
1698  
1699  
1700  
1701  
1702  
1703  
1704  
1705  
1706  
1707  
1708  
1709  
1710  
1711  
1712  
1713  
1714  
1715  
1716  
1717  
1718  
1719  
1720  
1721  
1722  
1723  
1724  
1725  
1726  
1727  
1728  
1729  
1730  
1731  
1732  
1733  
1734  
1735  
1736  
1737  
1738  
1739  
1740  
1741  
1742  
1743  
1744  
1745  
1746  
1747  
1748  
1749  
1750  
1751  
1752  
1753  
1754  
1755  
1756  
1757  
1758  
1759  
1760  
1761  
1762  
1763  
1764  
1765  
1766  
1767  
1768  
1769  
1770  
1771  
1772  
1773  
1774  
1775  
1776  
1777  
1778  
1779  
1780  
1781  
1782  
1783  
1784  
1785  
1786  
1787  
1788  
1789  
1790  
1791  
1792  
1793  
1794  
1795  
1796  
1797  
1798  
1799  
1800  
1801  
1802  
1803  
1804  
1805  
1806  
1807  
1808  
1809  
1810  
1811  
1812  
1813  
1814  
1815  
1816  
1817  
1818  
1819  
1820  
1821  
1822  
1823  
1824  
1825  
1826  
1827  
1828  
1829  
1830  
1831  
1832  
1833  
1834  
1835  
1836  
1837  
1838  
1839  
1840  
1841  
1842  
1843  
1844  
1845  
1846  
1847  
1848  
1849  
1850  
1851  
1852  
1853  
1854  
1855  
1856  
1857  
1858  
1859  
1860  
1861  
1862  
1863  
1864  
1865  
1866  
1867  
1868  
1869  
1870  
1871  
1872  
1873  
1874  
1875  
1876  
1877  
1878  
1879  
1880  
1881  
1882  
1883  
1884  
1885  
1886  
1887  
1888  
1889  
1890  
1891  
1892  
1893  
1894  
1895  
1896  
1897  
1898  
1899  
1900  
1901  
1902  
1903  
1904  
1905  
1906  
1907  
1908  
1909  
1910  
1911  
1912  
1913  
1914  
1915  
1916  
1917  
1918  
1919  
1920  
1921  
1922  
1923  
1924  
1925  
1926  
1927  
1928  
1929  
1930  
1931  
1932  
1933  
1934  
1935  
1936  
1937  
1938  
1939  
1940  
1941  
1942  
1943  
1944  
1945  
1946  
1947  
1948  
1949  
1950  
1951  
1952  
1953  
1954  
1955  
1956  
1957  
1958  
1959  
1960  
1961  
1962  
1963  
1964  
1965  
1966  
1967  
1968  
1969  
1970  
1971  
1972  
1973  
1974  
1975  
1976  
1977  
1978  
1979  
1980  
1981  
1982  
1983  
1984  
1985  
1986  
1987  
1988  
1989  
1990  
1991  
1992  
1993  
1994  
1995  
1996  
1997  
1998  
1999  
2000  
2001  
2002  
2003  
2004  
2005  
2006  
2007  
2008  
2009  
2010  
2011  
2012  
2013  
2014  
2015  
2016  
2017  
2018  
2019  
2020  
2021  
2022  
2023  
2024  
2025  
2026  
2027  
2028  
2029  
2030  
2031  
2032  
2033  
2034  
2035  
2036  
2037  
2038  
2039  
2040  
2041  
2042  
2043  
2044  
2045  
2046  
2047  
2048  
2049  
2050  
2051  
2052  
2053  
2054  
2055  
2056  
2057  
2058  
2059  
2060  
2061  
2062  
2063  
2064  
2065  
2066  
2067  
2068  
2069  
2070  
2071  
2072  
2073  
2074  
2075  
2076  
2077  
2078  
2079  
2080  
2081  
2082  
2083  
2084  
2085  
2086  
2087  
2088  
2089  
2090  
2091  
2092  
2093  
2094  
2095  
2096  
2097  
2098  
2099  
2100  
2101  
2102  
2103  
2104  
2105  
2106  
2107  
2108  
2109  
2110  
2111  
2112  
2113  
2114  
2115  
2116  
2117  
2118  
2119  
2120  
2121  
2122  
2123  
2124  
2125  
2126  
2127  
2128  
2129  
2130  
2131  
2132  
2133  
2134  
2135  
2136  
2137  
2138  
2139  
2140  
2141  
2142  
2143  
2144  
2145  
2146  
2147  
2148  
2149  
2150  
2151  
2152  
2153  
2154  
2155  
2156  
2157  
2158  
2159  
2160  
2161  
2162  
2163  
2164  
2165  
2166  
2167  
2168  
2169  
2170  
2171  
2172  
2173  
2174  
2175  
2176  
2177  
2178  
2179  
2180  
2181  
2182  
2183  
2184  
2185  
2186  
2187  
2188  
2189  
2190  
2191  
2192  
2193  
2194  
2195  
219

Sub B1

## BACKGROUND OF THE INVENTION

The United States Government may own certain rights in this invention under DARPA Grant No.: N65236-98-1-5413 and MURI Grant No.: DAAD19-99-1-0207.

5 This patent claims priority from provisional patent application Serial No.: 60/212,097, entitled "Aptazymes for Genetic Regulatory Circuits", by Andrew D. Ellington, et al., filed June 15, 2000.

10 Ribozymes or RNA enzymes are oligonucleotides of RNA that can act like enzymes by catalyzing the cleavage of RNA molecules. Generally, ribozymes have the ability to behave like an endoribonucleases. The location of the cleavage site is highly sequence specific, approaching the sequence specificity of DNA restriction endonucleases. By varying  
15 conditions, ribozymes can also act as polymerases or dephosphorylases.

20 Ribozymes were first described in connection with *Tetrahymena thermophila*. The *Tetrahymena* rRNA was shown to contain an intervening sequence (IVS) capable of excising itself out of a large ribosomal RNA precursor. The IVS is a catalytic RNA molecule that mediates self-splicing out of a precursor, whereupon it converts itself into a circular form.

The Tetrahymena IVS is more commonly known now as the Group I Intron.

Regulatable ribozymes have been described, wherein the activity of the ribozyme is regulated by a ligand binding moiety. Upon binding the ligand, the ribozyme activity on a target RNA is changed. The ligand-binding portion is an RNA sequence capable of binding a ligand such as an organic or inorganic molecule, or even a prodrug. The regulatable ribozymes described to date target bind, e.g., a first target sequence and the enzymatic activity is brought to bear on a separate RNA molecule for cleavage.

Ribozymes provide a mechanism to control the expression of genes in vitro and in vivo and therefore the expression of genes in patients. As such, ribozyme-based gene therapies using ribozymes may provide great medical benefits.

## SUMMARY OF THE INVENTION

5 The present invention provides an allosterically regulatable ribozyme or aptazyme that has the advantage of specific gene recognition with modulation of the enzymatic activity of the gene product typically exploited by pharmaceuticals. The aptazymes of the present invention are, therefore, allosteric ribozymes in that their activity is under the allosteric control of a second portion of the ribozyme. Just as allosteric protein enzymes undergo a change in their kinetic parameters or of their enzymatic activity in response to interactions with an effector molecule, the catalytic abilities of the regulatable aptazymes may similarly be modulated by an allosteric effector(s). Thus, the present invention is directed to allosterically regulatable aptazymes that transduce molecular recognition into catalysis.

10  
15  
20 The present invention includes an allosterically regulatable aptazyme construct that is inserted into a gene of interest, e.g., a gene targeting expression vector. The regulatable aptazyme sequence provides gene specific recognition as well as modulation of the aptazyme's kinetic parameters. The kinetic parameters of the regulatable aptazyme vary in response to an allosteric effector molecule. Specifically, in the presence of the allosteric effector, the

aptazyme splices itself out of the gene in response to the effector molecule to regulate expression of the gene. An important feature of the present invention is that the regulatable aptazymes disclosed herein only require recognition rather than actual binding.

A key distinction with known systems is that the regulatory domains of the regulatable aptazymes of the present invention may bind targets, but they are engineered and selected without the necessity of knowing anything about their binding target. In the present invention it is the allosteric interaction of an effector molecule with the regulatory domain that transduces interactions into catalysis. Therefore, binding is a concomitant but secondary function of such interactions; that is, the allosteric ribozymes disclosed herein may bind the effector or the target very poorly, but upon their interaction, a synergistic effect is found that could not be detected by screening for each characteristic alone.

It is yet another embodiment of the present invention that the effector molecule does not only (or will not) produce a conformational change, but rather will add essential catalytic sites (e.g., residues) for a reaction. That is, both the effector molecule and the regulatable aptazyme contribute a portion of the active site of the ribozyme. For

example, using the method of the present invention a ribozyme and an effector molecule that would be unable to bind and/or perform an enzymatic function independently, may be isolated that act synergistically. As such, an regulatable aptazyme that contributes a guanosine and an adenosine and an effector molecule that contributes a histidine from a protein effector form a synergistic effector-aptazyme complex that is regulatable based on the presence and concentration of the effector or the aptazyme. Using the methods disclosed herein it is possible to identify a chimeric effector:aptazyme (e.g., a protein:RNA complex) active site that would lead to catalysis.

The present invention also includes an aptazyme construct with a regulatable aptamer oligonucleotide sequence having a regulatory domain, such that the kinetic parameters of the aptazyme on a target gene vary in response to the interaction of an allosteric effector molecule with the regulatory domain.

The regulatable aptazyme may be used for gene expression, up regulation (increasing production of the gene product) or down regulation (decreasing the production of the gene product). The construct of one embodiment of the present invention provides a DNA oligonucleotide coding for an aptazyme domain so that the DNA can be transcribed to RNA (e.g., mRNA), where the RNA contains a self-splicing aptazyme

domain that can be activated in the presence of an effector molecule. The advantages of the aptazyme technology of the present invention include the ability to continually modulate gene expression with a high degree of sensitivity without additional gene therapy interventions.

Aptazymes are more robust than allosteric protein enzymes in several ways: (1) they can be selected in vitro, which facilitates the engineering of particular constructs; (2) the levels of catalytic modulation are much greater for aptazymes than for protein enzymes; and (3) since aptazymes are nucleic acids, they can potentially interact with the genetic machinery in ways that protein molecules may not.

The regulatable aptazymes of the present invention may also be expressed inside cells. The regulatable aptazymes of the present invention that are expressed inside a cell are not only responsive to a given effector molecule, but are also able to participate in genetic regulation or responsiveness. In particular, self-splicing introns can splice themselves out of genes in response to exogenous or endogenous effector molecules.

For example, a gene can be activated or repressed in response to an exogenously introduced allosteric effector (drug) for gene therapy. In fact, at least part of the

utility of the present invention is for use in the identification, isolation and enhancement of allosteric effectors and of the allosterically regulatable aptazymes with which they interact. Similarly, it is possible to activate or repress a reporter gene (e.g., luciferase) containing an engineered intron in response to an endogenous activator. In this way, luciferase-engineered intron constructs may be used to monitor intracellular levels of proteins or small molecules such as cyclic AMP. Such applications may be used for high-throughput screening. If a particular intracellular signal (e.g., the production of a tumor repressor) was desired, compound libraries are screened for pharmacophores that induce the signal (the tumor repressor), which in turn activates the intron and leads to the production of a detectable signal (e.g., expression of luciferase). Thus, the information desired is changed or morphed into the detection of glowing cells.

One important feature of using regulatable aptazymes for gene therapies is that regulated introns are a generalizable means of controlling gene expression, for any of a variety of genes, since the introns could be inserted into and be engineered to accommodate virtually any gene. Moreover, since the regulatable aptazymes may be engineered to respond to any



of a variety of effectors, the characteristics of the effector (oral availability, synthetic accessibility, pharmacokinetic properties) may be chosen in advance. The drug is chosen prior to engineering the target of the drug. In part because of these extraordinary capabilities, aptazymes provide perhaps a powerful route to medically successful and practical gene therapies. Drugs may be given throughout the treatment (or lifetime) of a patient who had undergone a single initial gene therapy. By making the gene therapy regulatable, the amount of a gene product may be easily increased or decreased in different individuals at different times during the treatment by increasing or decreasing the doses of effector molecules.

The present invention includes a method for providing a regulatable aptazyme construct having an aptamer oligonucleotide sequence with a regulatory domain. A characteristic of the regulatable aptazyme construct of the present method is that the kinetic parameters of the aptazyme vary in response to an effector molecule. In particular, the kinetic parameters of the aptazyme on a target gene vary in response to the interaction of an allosteric effector molecule with the regulatory domain. For example, the aptazyme splices itself out of a gene in response to the effector molecule

interacting with the regulatory domain of the aptazyme to regulate expression of the target gene.

The present method also includes transforming a cell with the regulatable aptazyme construct so that the aptazyme construct is inserted into a gene of interest. An effector molecule is provided to activate the aptazyme so that administering to the cell an effective amount of the allosteric effector molecule induces the aptazyme to splice itself out of the gene to regulate expression of the gene.

The method of the present invention contemplates that the aptazyme construct may be within a plasmid. The method then further includes transforming the cell with the aptazyme construct containing plasmid. The method of the present invention also includes ligating the regulatable aptazyme construct into a vector and transforming the cell with the vector. Additionally, the method of the present invention contemplates that the regulatable aptazyme construct may be amplified by polymerase chain reaction. Finally, the method contemplates that the regulatable aptamer oligonucleotide sequence of the construct may include RNA nucleotides, so that the method further includes reverse transcription of the RNA using reverse transcriptase to produce an DNA aptamer complementary to the RNA template.

The invention also includes the automation of *in vitro* selection, and a mechanized system that executes both common and modified *in vitro* selection procedures. Automation facilitates the execution of this procedure, accomplishing in hours-to-days what once necessitated weeks-to-months. Additionally, the mechanized system attends to other technical obstacles not addressed in 'common' *in vitro* selection procedure (e.g, specialized robotic manipulation to avoid cross-contamination).

The automation methods are generalizable to a number of different types of selections, including selections with DNA or modified RNA, selections for ribozymes and selections for phage-displayed or cell-surface displayed proteins.

Automating selection greatly diminishes human error in the actual pipetting and biological manipulations. While programming the robot is often not a trivial task, and can be time consuming, automated selection is far faster and more efficient than manual selection. The scientist's time is thus put to better use preparing samples and analyzing data, rather than performing the actual selection. Additionally, automated selection may include real-time monitoring methods (e.g., molecular beacons, TaqMan®) into the selection procedure and software that can make intelligent decisions based on real-time monitoring.

5

10

15

20

Page B3

Aptazymes, or allosterically activated ribozymes, have been developed that are activated cyclic nucleotide monophosphates as well as other small molecules like theophylline. In addition to aptazymes activated by small molecules, the present invention also includes allosteric ribozymes that are activated by protein cofactors. Indeed, there are natural ribozymes that are extremely dependent on proteins for their activity.

One embodiment of the present invention involves the in vitro selection of ribozymes that are allosterically activated by proteins. A selection scheme for ribozymes that are dependent on protein cofactors has been developed. A novel class of nucleoprotein enzymes has been identified where the enzymes are even more dependent on Cyt18 for their activity than is the natural Group I intron. Aptazymes may be allosterically regulated by any of a vast number of proteins.

Other methods for aptazyme development using small molecule ligands have proven successful. In particular, it has been possible to add aptamer moieties to ribozymes, without selection, and achieve activation in the presence of ligand (like ATP or theophylline). However, this has proved difficult in the case of protein ligands. One of the unique features of the present selection protocol relative to others

that have previously been published is that the present invention randomizes not only a domain that is pendant on the catalytic core, but a portion of the catalytic core itself. Thus, the selection for protein-dependence not only yields  
5 species which bind to ancillary regions of the ribozyme, but that likely help organize the catalytic core of the ribozyme.

It should be noted that the method is not limited to RNA pools, but could also encompass DNA pools or modified RNA pools. The method is not limited to ligases, but could also  
10 encompass other ribozyme classes. The method is not limited to protein-induced conformational changes, but could also take into account 'chimeric' catalysts in which residues essential for chemical reactivity were provided by both the nucleic acid and the protein, in concert.

This invention allows the selection of protein-dependent aptazymes, which are extremely novel reagents that can be  
15 useful in a variety of applications. For example, protein-dependent aptazymes can be used (1) in substrates for the acquisition of data about whole proteomes, (2) as in vitro diagnostic reagents to detect proteins specific to disease  
20 states, such as prostate-specific antigen (PSA) or viral proteins, (3) as sentinels for the detection of biological

warfare agents, or (4) as regulatory elements in gene therapies, as described herein.

Initially, many protein targets may prove refractive to selection. However, many derivatives of the base method can be developed, to deal with novel targets or target classes.

It has been shown that ribozyme catalysis can be modulated by allosteric effectors. In yet another embodiment of the present invention, these allosteric ribozymes, also referred to as aptazymes, are displayed in arrays to be used for monitoring the presence of various molecules, be they inorganic or organic (e.g., metabolites or proteins).

For example, aptazymes are anchored to a substrate, such as wells in a multi-well plate, and different ribozyme ligases are covalently immobilized on beads in the wells. Mixtures of analytes and fluorescently tagged substrates are added to each well. Where cognate effectors are present, the aptazymes will covalently attach the fluorescent tags to themselves. Where aptazymes have not been activated by effectors, the tagged substrates are washed out of the well. After reaction and washing, the presence and amounts of co-immobilized fluorescent tags are indicative of amounts of ligands that were present during the reaction.

In this embodiment of the invention, the reporter may be a fluorescent tag, but it may also be an enzyme, a magnetic particle, or any number of detectible particles. Additionally, the ribozymes could be immobilized on beads, but they could also be directly attached to a solid support via covalent bonds.

One advantage of this embodiment is that covalent immobilization of reporters (as opposed to non covalent immobilization, as in ELISA™ assays) allows stringent wash steps to be employed. Additionally, ribozyme ligases have the unique property of being able to transduce effectors into templates that can be amplified, affording an additional boost in signal prior to detection.

Nucleic acids are generally less robust than antibodies. However, modified nucleotides may be introduced into the aptazymes that substantially stabilize them from degradation in environments such as sera or urine. Similarly, antibodies generally have higher affinities for analytes than do aptamers, and be inference aptazymes. However, the analytical methods of the present invention do not rely on binding per se, but only on transient interactions. The present invention requires mere recognition rather than actual binding, providing a potential advantage of aptazymes over antibodies.





## BRIEF DESCRIPTION OF THE DRAWINGS

For a more complete understanding of the features and advantages of the present invention, reference is now made to the detailed description of the invention along with the accompanying figures in which corresponding numerals in different figures refer to corresponding parts and in which:

Figure 1 is a depiction of the secondary structure of the Group 1 theophylline-dependent (td) intron of bacteriophage T4 (wild type);

Figure 2a is a photograph of a gel showing activation of the GpITh1P6.131 aptamer construct, together with a graphical representation of the gel, of one embodiment of the present invention;

Figure 2b is a photograph of a gel showing activation of GpITh2P6.133 aptamer construct, together with a graphical representation of the gel of one embodiment of the present invention.

Figure 3 is a schematic depiction of an in vivo assay system for group I introns of one embodiment of the present invention.

Figure 4a depicts a portion of the P6 region of the Group I ribozyme joined to the anti-theophylline aptamer by a

short randomized region to generate a pool of aptazymes of the present invention.

Figure 4b is a schematic depiction of a selection protocol for the Group I P6 Aptazyme Pool of Fig. 4a.

5        Figure 5 is a diagram of one embodiment of the present invention depicting exogenous or endogenous activation of Group I intron splicing;

10       Figure 6 is a diagram of another embodiment of the present invention depicting a strategy for screening libraries of exogenous activators;

Figure 7 is a diagram of an alternative embodiment of the present invention for screening libraries of exogenous activators;

15       Figure 8 is a diagram of yet another alternative embodiment of the present invention for screening libraries of exogenous activators;

Figure 9 is a diagram of an embodiment of the present invention for screening for endogenous activators;

20       Figure 10 is a diagram of an alternative to the embodiment of Fig. 9 of the present invention to screen for endogenous activators;

Figure 11 is a diagram of another embodiment of the present invention to screen for compounds that perturb cellular metabolism;

Figure 12 is a diagram of a further embodiment of the present invention that provides a non-invasive readout of metabolic states;

Figure 13 is a diagram of yet a further embodiment of the present invention wherein endogenous suppressors provide a non-invasive readout of multiple metabolic states;

Figure 14 is a schematic depiction of an example of a worksurface for automatic selection procedures of one embodiment of the invention;

Figure 15a is an illustration of the LI ligase aptazyme construct of one embodiment of the present invention;

Figure 15b is an illustration of a modified LI ligase aptazyme construct of Figure 15a of the present invention;

Figure 15c is a schematic diagram of a selection protocol of one embodiment of the present invention; and

Figure 16 is a schematic diagram of a method to anchor an aptazyme construct of the present invention to a solid support in one embodiment of the present invention.

## DETAILED DESCRIPTION OF THE INVENTION

While the making and using of various embodiments of the present invention are discussed in detail below, it should be appreciated that the present invention provides many applicable inventive concepts that may be embodied in a wide variety of specific contexts. The specific embodiments discussed herein are merely illustrative of specific ways to make and use the invention and do not delimit the scope of the invention.

### Definitions

As used herein, the term "regulatable aptazyme" means an allosteric ribozyme. The kinetic parameters of the ribozyme may be varied in response to the amount of an allosteric effector molecule. Just as allosteric protein enzymes undergo a change in their kinetic parameters or of their enzymatic activity in response to interactions with an effector molecule, the catalytic abilities of regulatable aptazymes can similarly be modulated by effectors. Regulatable aptazymes transduce molecular recognition into catalysis upon interaction with an allosteric effector molecule that binds an effector portion of the regulatable aptazyme. Specifically, in the presence of the effector, the aptazyme splices itself

out of a gene in response to the effector molecule to regulate expression of the gene.

As used herein, the term "aptamer" refers to an oligonucleotide having aptazyme activity.

5 As used herein, the term "allosteric effector" or "allosteric effector molecule" means a substance that allosterically changes the kinetic parameters or catalytic activity of an aptazyme, and in particular a substance that activates self-splicing of an aptazyme.

10 As used herein, the term "kinetic parameters" refers to catalytic activity. Changes in the kinetic parameters of a catalytic ribozyme produce changes in the catalytic activity of the ribozyme such as a change in the rate of reaction or a change in substrate specificity. For example, self-splicing  
15 of an aptazyme out of a gene environment may result from a change in the kinetic parameters of the aptazyme.

As used herein, the term "catalytic" or "catalytic activity" refers to the ability of a substance to affect a change in itself or of a substrate under permissive  
20 conditions.

As used herein, the term "protein-protein complex" or "protein complex" refers to an association of more than one protein. The proteins of the complex may be associated by a

variety of means, or by any combination of means, including but not limited to functional, stereochemical, conformational, biochemical, or electrostatic association. It is intended that the term encompass associations of any number of proteins.

As used herein the terms "protein", "polypeptide" or "peptide" refer to compounds comprising amino acids joined via peptide bonds and are used interchangeably.

As used herein, the term "endogenous" refers to a substance the source of which is from within a cell. Endogenous substances are produced by the metabolic activity of a cell. Endogenous substances, however, may nevertheless be produced as a result of manipulation of cellular metabolism to, for example, make the cell express the gene encoding the substance.

As used herein, the term "exogenous" refers to a substance the source of which is external to a cell. An exogenous substance may nevertheless be internalized by a cell by any one of a variety of metabolic or induced means known to those skilled in the art.

As used herein, the term "gene" means the coding region of a deoxyribonucleotide sequence encoding the amino acid sequence of a protein. The term includes sequences located

adjacent to the coding region on both the 5', and 3', ends such that the deoxyribonucleotide sequence corresponds to the length of the full-length mRNA for the protein. The term "gene" encompasses both cDNA and genomic forms of a gene. A genomic form or clone of a gene contains the coding region interrupted with non-coding sequences termed "introns" or "intervening regions" or "intervening sequences." Introns are segments of a gene which are transcribed into nuclear RNA (hnRNA); introns may contain regulatory elements such as enhancers. Introns are removed, excised or "spliced out" from the nuclear or primary transcript; introns therefore are absent in the messenger RNA (mRNA) transcript. The mRNA functions during translation to specify the sequence or order of amino acids in a nascent polypeptide.

In addition to containing introns, genomic forms of a gene may also include sequences located on both the 5' and 3' end of the sequences which are present on the RNA transcript. These sequences are referred to as "flanking" sequences or regions (these flanking sequences are located 5' or 3' to the non-translated sequences present on the mRNA transcript). The 5' flanking region may contain regulatory sequences such as promoters and enhancers which control or influence the transcription of the gene. The 3' flanking region may contain

87 Contd

sequences which direct the ~~termination~~ of transcription, post-transcriptional cleavage and polyadenylation.

DNA molecules are said to have "5' ends" and "3' ends" because mononucleotides are reacted to make oligonucleotides in a manner such that the 5' phosphate of one mononucleotide pentose ring is attached to the 3' oxygen of its neighbor in one direction via a phosphodiester linkage. Therefore, an end of an oligonucleotides referred to as the "5' end" if its 5' phosphate is not linked to the 3' oxygen of a mononucleotide pentose ring and as the "3' end" if its 3' oxygen is not linked to a 5' phosphate of a subsequent mononucleotide pentose ring. As used herein, a nucleic acid sequence, even if internal to a larger oligonucleotide, also may be said to have 5' and 3' ends. In either a linear or circular DNA molecule, discrete elements are referred to as being "upstream" or 5' of the "downstream" or 3' elements. This terminology reflects the fact that transcription proceeds in a 5' to 3' fashion along the DNA strand.

The term "gene of interest" as used here refers to a gene, the function and/or expression of which is desired to be investigated, or the expression of which is desired to be regulated, by the present invention. In the present disclosure, the *td* gene of the T4 bacteriophage is an example



of a gene of interest and is described herein to illustrate the invention. The present invention may be useful in regard to any gene of any organism, whether of a prokaryotic or eukaryotic organism.

5           The term "hybridize" as used herein, refers to any process by which a strand of nucleic acid binds with a complementary strand through base pairing. Hybridization and the strength of hybridization (i.e., the strength of the association between the nucleic acid strands) is impacted by  
10 such factors as the degree of complementary between the nucleic acids, stringency of the conditions involved, the melting temperature of the formed hybrid, and the G:C (or U:C for RNA) ratio within the nucleic acids.

15           The terms "complementary" or "complementarity" as used herein, refer to the natural binding of polynucleotides under permissive salt and temperature conditions by base-pairing. For example, for the sequence "A-G-T" binds to the complementary sequence "T-C-A". Complementarity between two single-stranded molecules may be partial, in which only some  
20 of the nucleic acids bind, or it may be complete when total complementarity exists between the single stranded molecules. The degree of complementarity between nucleic acid strands has significant effects on the efficiency and strength of

hybridization between nucleic acid strands. This is of particular importance in amplification reactions, which depend upon binding between nucleic acids strands.

The term "homology," as used herein, refers to a degree of complementarity. There may be partial homology or complete homology (*i.e.*, identity). A partially complementary sequence is one that at least partially inhibits an identical sequence from hybridizing to a target nucleic acid; it is referred to using the functional term "substantially homologous." The inhibition of hybridization of the completely complementary sequence to the target sequence may be examined using a hybridization assay (Southern or Northern blot, solution hybridization and the like) under conditions of low stringency. A substantially homologous sequence or probe will compete for and inhibit the binding (*i.e.*, the hybridization) of a completely homologous sequence or probe to the target sequence under conditions of low stringency. This is not to say that conditions of low stringency are such that non-specific binding is permitted; low stringency conditions require that the binding of two sequences to one another be a specific (*i.e.*, selective) interaction. The absence of non-specific binding may be tested by the use of a second target sequence which lacks even a partial degree of complementarity

(e.g., less than about 30% identity); in the absence of non-specific binding, the probe will not hybridize to the second non-complementary target sequence. When used in reference to a single-stranded nucleic acid sequence, the term "substantially homologous" refers to any probe which can hybridize (i.e., it is the complement of) the single-stranded nucleic acid sequence under conditions of low stringency as described.

As known in the art, numerous equivalent conditions may be employed to comprise either low or high stringency conditions. Factors such as the length and nature (DNA, RNA, base composition) of the sequence, nature of the target (DNA, RNA, base composition, presence in solution or immobilization, etc.), and the concentration of the salts and other components (e.g., the presence or absence of formamide, dextran sulfate and/or polyethylene glycol) are considered and the hybridization solution may be varied to generate conditions of either low or high stringency different from, but equivalent to, the above listed conditions.

As used herein the term "stringency" is used in reference to the conditions of temperature, ionic strength, and the presence of other compounds such as organic solvents, under which nucleic acid hybridizations are conducted. With "high

stringency" conditions, nucleic acid base pairing will occur only between nucleic acid fragments that have a high frequency of complementary base sequences. Thus, conditions of "weak" or "low" stringency are often required with nucleic acids that are derived from organisms that are genetically diverse, as the frequency of complementary sequences is usually less.

Low stringency conditions comprise conditions equivalent to binding or hybridization at 42°C in a solution consisting of 5X SSPE (43.8 g/l NaCl, 6.9 g/l NaH<sub>2</sub>PO<sub>4</sub>•H<sub>2</sub>O and 1.85 g/l EDTA, pH adjusted to 7.4 with NaOH), 0.1% SDS, 5X Denhardt's reagent (50X Denhardt's contains per 500 ml: 5 g Ficoll (Type 400, Pharmacia), 5 g BSA [Fraction V; Sigma]) and 100 µg/ml denatured salmon sperm DNA) followed by washing in a solution comprising 5X SSPE, 0.1% SDS at 42°C when a probe of about 500 nucleotides in length is employed.

Numerous equivalent conditions may be employed to comprise low stringency conditions; factors such as the length and nature (DNA, RNA, base composition) of the probe and nature of the target (DNA, RNA, base composition, present in solution or immobilized, etc.) and the concentration of the salts and other components (e.g., the presence or absence of formamide, dextran sulfate, polyethylene glycol) are considered and the hybridization solution may be varied to

generate conditions of low stringency hybridization different from, but equivalent to, the above listed conditions. In addition, the art knows conditions which promote hybridization under conditions of high stringency (e.g., increasing the temperature of the hybridization and/or wash steps, the use of formamide in the hybridization solution, etc.).

The term "antisense," as used herein, refers to nucleotide sequences that are complementary to a specific DNA or RNA sequence. The term "antisense strand" is used in reference to a nucleic acid strand that is complementary to the "sense" strand. Antisense molecules may be produced by any method, including synthesis by ligating the gene(s) of interest in a reverse orientation to a viral promoter which permits the synthesis of a complementary strand. Once introduced into a cell, the transcribed strand combines with natural sequences produced by the cell to form duplexes. These duplexes then block either the further transcription or translation. In this manner, mutant phenotypes may also be generated. The designation "negative" is sometimes used in reference to the antisense strand, and "positive" is sometimes used in reference to the sense strand.

The term also is used in reference to RNA sequences that are complementary to a specific RNA sequence (e.g., mRNA).

Included within this definition are antisense RNA ("asRNA") molecules involved in genetic regulation by bacteria. Antisense RNA may be produced by any method, including synthesis by splicing the gene(s) of interest in a reverse orientation to a viral promoter which permits the synthesis of a coding strand. Once introduced into an embryo, this transcribed strand combines with natural mRNA produced by the embryo to form duplexes. These duplexes then block either the further transcription of the mRNA or its translation. In this manner, mutant phenotypes may be generated. The term "antisense strand" is used in reference to a nucleic acid strand that is complementary to the "sense" strand. The designation. (-) (i.e., "negative") is sometimes used in reference to the antisense strand with the designation (+) sometimes used in reference to the sense (i.e., "positive") strand.

A gene may produce multiple RNA species which are generated by differential splicing of the primary RNA transcript. cDNAs that are splice variants of the same gene will contain regions of sequence identity or complete homology (representing the presence of the same exon or portion of the same exon on both cDNAs) and regions of complete non-identity (for example, representing the presence of exon "A" on cDNA I

wherein cDNA 2 contains exon "B" instead). Because the two cDNAs contain regions of sequence identity they will both hybridize to a probe derived from the entire gene or portions of the gene containing sequences found on both cDNAs; the two splice variants are therefore substantially homologous to such a probe and to each other.

"Transformation," as defined herein, describes a process by which exogenous DNA enters and changes a recipient cell. It may occur under natural or artificial conditions using various methods well known in the art. Transformation may rely on any known method for the insertion of foreign nucleic acid sequences into a prokaryotic or eukaryotic host cell. The method is selected based on the host cell being transformed and may include, but is not limited to, viral infection, electroporation, lipofection, and particle bombardment. Such "transformed" cells include stably transformed cells in which the inserted DNA is capable of replication either as an autonomously replicating plasmid or as part of the host chromosome. The term "transfection" as used herein refers to the introduction of foreign DNA into eukaryotic cells.

Transfection may be accomplished by a variety of means known to the art including, e.g., calcium phosphate-DNA co-

precipitation, DEAE-dextran-mediated transfection, polybrene-mediated transfection, electroporation, microinjection, liposome fusion, lipofection, protoplast fusion, retroviral infection, and biolistics. Thus, the term "stable transfection" or "stably transfected" refers to the introduction and integration of foreign DNA into the genome of the transfected cell. The term "stable transfectant" refers to a cell which has stably integrated foreign DNA into the genomic DNA. The term also encompasses cells which transiently express the inserted DNA or RNA for limited periods of time. Thus, the term "transient transfection" or "transiently transfected" refers to the introduction of foreign DNA into a cell where the foreign DNA fails to integrate into the genome of the transfected cell. The foreign DNA persists in the nucleus of the transfected cell for several days. During this time the foreign DNA is subject to the regulatory controls that govern the expression of endogenous genes in the chromosomes. The term "transient transfectant" refers to cells which have taken up foreign DNA but have failed to integrate this DNA.

As used herein, the term "selectable marker" refers to the use of a gene that encodes an enzymatic activity and which confers the ability to grow in medium lacking what would

Sub B8



Sub B8  
Contd

otherwise be an essential nutrient (e.g., the HIS3 gene in yeast cells); in addition, a selectable marker may confer resistance to an antibiotic or drug upon the cell in which the selectable marker is expressed. A review of the use of selectable markers in mammalian cell lines is provided in Sambrook, J. et. al., *Molecular Cloning: A Laboratory Manual*, 2nd ed., Cold Spring Harbor Laboratory Press, New York (1989) pp.16.9-16.15.

As used herein, the term "reporter gene" refers to a gene that is expressed in a cell upon satisfaction of one or more contingencies and which, upon expression, confers a detectable phenotype to the cell to indicate that the contingencies for expression have been satisfied. For example, the gene for Luciferase confers a luminescent phenotype to a cell when the gene is expressed inside the cell. In the present invention, the gene for Luciferase may be used as a reporter gene such that the gene is only expressed upon the splicing out of an intron in response to an effector. Those cells in which the effector activates splicing of the intron will express Luciferase and will glow.

As used herein, the term "vector" is used in reference to nucleic acid molecules that transfer DNA segment(s) from one

cell to another. The term "vehicle" is sometimes used interchangeably with "vector."

The term "vector" as used herein also includes expression vectors in reference to a recombinant DNA molecule containing a desired coding sequence and appropriate nucleic acid sequences necessary for the expression of the operably linked coding sequence in a particular host organism. Nucleic acid sequences necessary for expression in prokaryotes usually include a promoter, an operator (optional), and a ribosome binding site, often along with other sequences. Eukaryotic cells are known to utilize promoters, enhancers, and termination and polyadenylation signals.

As used herein, the term "amplify", when used in reference to nucleic acids refers to the production of a large number of copies of a nucleic acid sequence by any method known in the art. Amplification is a special case of nucleic acid replication involving template specificity. Template specificity is frequently described in terms of "target" specificity. Target sequences are "targets" in the sense that they are sought to be sorted out from other nucleic acid. Amplification techniques have been designed primarily for this sorting out.

As used herein, the term "primer" refers to an oligonucleotide, whether occurring naturally as in a purified restriction digest or produced synthetically, which is capable of acting as a point of initiation of synthesis when placed under conditions in which synthesis of a primer extension product which is complementary to a nucleic acid strand is induced, (i.e., in the presence of nucleotides and an inducing agent such as DNA polymerase and at a suitable temperature and pH). The primer may be single stranded for maximum efficiency in amplification but may alternatively be double stranded. If double stranded, the primer is first treated to separate its strands before being used to prepare extension products. The primer must be sufficiently long to prime the synthesis of extension products in the presence of the inducing agent. The exact lengths of the primers will depend on many factors, including temperature, source of primer and the use of the method.

As used herein, the term "probe" refers to an oligonucleotide (i.e., a sequence of nucleotides), whether occurring naturally as in a purified restriction digest or produced synthetically, recombinantly or by PCR amplification, which is capable of hybridizing to another oligonucleotide of interest. A probe may be single-stranded or double-stranded.

Sub B9  
Contd.

Probes are useful in the detection, identification and isolation of particular gene sequences. It is contemplated that any probe used in the present invention will be labeled with any "reporter molecule," so that is detectable in any detection system, including, but not limited to enzyme (e.g. ELISA, as well as enzyme-based histochemical assays), fluorescent, radioactive, and luminescent systems. It is not intended that the present invention be limited to any particular detection system or label.

10

Sub B9

As used herein, the term "target" when used in reference to the polymerase chain reaction, refers to the region of nucleic acid bounded by the primers used for polymerase chain reaction. Thus, the "target" is sought to be sorted out from other nucleic acid sequences. A "segment" is defined as a region of nucleic acid within the target sequence.

15

20

As used herein, the term "polymerase chain reaction" ("PCR") refers to the method of K.B. Mullis U.S. Patent Nos. 4,683,195, 4,683,202, and 4,965,188, hereby incorporated by reference, which describe a method for increasing the concentration of a segment of a target sequence in a mixture of genomic DNA without cloning or purification. This process for amplifying the target sequence consists of introducing a large excess of two oligonucleotide primers to the DNA mixture

containing the desired target sequence, followed by a precise sequence of thermal cycling in the presence of a DNA polymerase. The two primers are complementary to their respective strands of the double stranded target sequence.

5           To effect amplification, the mixture is denatured and the primers then annealed to their complementary sequences within the target molecule. Following annealing, the primers are extended with a polymerase so as to form a new pair of complementary strands. The steps of denaturation, primer annealing and polymerase extension can be repeated many times (i.e., denaturation, annealing and extension constitute one "cycle"; there can be numerous "cycles") to obtain a high concentration of an amplified segment of the desired target sequence. The length of the amplified segment of the desired target sequence is determined by the relative positions of the primers with respect to each other, and therefore, this length is a controllable parameter. By virtue of the repeating aspect of the process, the method is referred to as the "polymerase chain reaction" (hereinafter "PCR"). Because the  
10  
15  
20 desired amplified segments of the target sequence become the predominant sequences (in terms of concentration) in the mixture, they are said to be "PCR amplified".

5 Sub B11

With PCR, it is possible to amplify a single copy of a specific target sequence in genomic DNA to a level detectable by several different methodologies (e.g., hybridization with a labeled probe; incorporation of biotinylated primers followed by avidin-enzyme conjugate detection; incorporation of <sup>32</sup>P-labeled deoxynucleotide triphosphates, such as DCTP or DATP, into the amplified segment). In addition to genomic DNA, any oligonucleotide sequence can be amplified with the appropriate set of primer molecules. In particular the amplified segments created by the PCR process itself are, themselves, efficient templates for subsequent PCR amplifications.

10

15 The invention is now described in detail with the use of the td intron from T4 bacteriophage (Fig. 1) for illustrative purposes. The description is not intended to limit the scope of the invention or the claims appended hereto.

20 Figure 1 depicts the secondary structure of the td intron from bacteriophage T4 (GenBank # M12742), wherein "td" means theophylline-dependent. The td intron was selected to illustrate the present invention because, among other things, mutational analysis has identified regions of this intron that can be engineered and modified. See Salvo, et al., Deletion-tolerance and trans-splicing of the bacteriophage T4 intron.

Analysis of the P6-L6a region. *J. Mol. Biol.* **211**, 537-549 (1990) and Salvo et al., The P2 element of the td intron is dispensable despite its normal role in splicing. *J. Mol. Biol.* **267**, 2845-2848 (1992). Thus, aptamer domains or pools may be engineered into the T4 intron.

The intron has been adapted to a prokaryotic in vivo screening protocol. M. Belfort, et al., Genetic delineation of functional components of the group I intron in the phage T4 td gene. *Cold Spring Harb Symp Quant Biol* **52**, 181-92 (1987). The present invention improves the Belfort, et al., protocol to assay drug dependence in vivo. The protocol may also be used to screen and select Group I aptazymes in vivo.

A final advantage of the T4 intron is that the intron can be rendered protein-dependent. C.A. Myers, et al., A tyrosyl-tRNA synthetase suppresses structural defects in the two major helical domains of the group I intron catalytic core. *J. Mol. Biol.* **262** 87--104 (1996). An anti-theophylline aptamer has been described by R.D. Jenison, et al., High-resolution molecular discrimination by RNA. *Science* **263**, 1425-1429 (1994).

In the present invention, the anti-theophylline aptamer was mounted in two locations in the td intron, shown by the shaded portions of Fig. 1. One location was at the termini of

P1 and the other location was within P6. The P1 constructs may enable ligand-dependent conformational changes that alter the conformation or register of the U:G base pair which is critical for splicing. The P6 region was selected because mutational analysis indicated that deletion of the P6 stem destabilizes the intron.

Referring now to Figs 2a and 2b, in the present invention P6 constructs were made so that Group I splicing was activated by the presence of theophylline in the range of approximately 9 to 19 fold over constructs grown in the absence of theophylline, as described in the examples below:

The following examples illustrate the present invention in the *td* gene system of T4. For a full understanding of the examples, refer to Figures 2a and 2b. The examples are provide for illustrative purposes and do not limit the scope of the present invention or the scope of the appended claims.

#### **Example 1: GpITH1P6**

##### **Engineering of Group I Aptazymes**

The first example illustrates how to make an aptazyme construct and demonstrates self-splicing of the aptazyme out of a gene in response to an effector molecule.

##### Construction of a regulatable aptazyme



Oligos GpIWt3.129: 5'-TAA TCT TAC CCC GGA ATT ATA TCC AGC  
TGC ATG TCA CCA TGC AGA GCA GAC TAT ATC TCC AAC TTG TTA AAG  
CAA GTT GTC TAT CGT TTC GAG TCA CTT GAC CCT ACT CCC CAA AGG  
GAT AGT CGT TAG-3' (SEQ ID NO: 1) and GpITh1P6.131: 5'-GCC TGA  
GTA TAA GGT GAC TTA TAC TTG TAA TCT ATC TAA ACG GGG AAC CTC  
TCT AGT AGA CAA TCC CGT GCT AAA TTA TAC CAG CAT CGT CTT GAT  
GCC CTT GGC AGA TAA ATG CCT AAC GAC TAT CCC TT-3' (SEQ ID NO:  
2) were annealed and extended in a 30  $\mu$ l reaction containing  
100 pmoles of each oligo, 250 mM Tris-HCl (pH 8.3), 40 mM  
MgCl<sub>2</sub>, 250 mM NaCl, 5 mM DTT, 0.2 mM each dNTP, 45 units of  
AMV reverse transcriptase (RT: Amersham Pharmacia Biotech,  
Inc., Piscataway, NJ) at 37° C for 30 minutes. The extension  
reaction was diluted 1 to 50 in H<sub>2</sub>O.

A PCR reaction containing 1  $\mu$ l of the extension dilution,  
500 mM KCl, 100 mM Tris-HCl (pH 9.0), 1% Triton® x-100, 15 mM  
MgCl<sub>2</sub>, 0.4  $\mu$ M of GpIWt1.75: 5'-GAT AAT ACG ACT CAC TAT AGG GAT  
CAA CGC TCA GTA GAT GTT TTC TTG GGT TAA TTG AGG CCT GAG TAT  
AAG GTG-3' (SEQ ID NO:3), 0.4  $\mu$ M of GpIWt4.89: 5'-CTT AGC TAC  
AAT ATG AAC TAA CGT AGC ATA TGA CGC AAT ATT AAA CGG TAG CAT  
TAT GTT CAG ATA AGG TCG TTA ATC TTA CCC CGG AA-3' (SEQ ID  
NO:4), 0.2 mM each dNTP and 1.5 units of Taq polymerase  
(Promega, Madison, WI) was thermocycled 20 times under the  
following regime: 94° C for 30 seconds, 45° C for 30 seconds,

Sub B 13  
Cont'd

72° C for 1 minute. The PCR reaction was precipitated in the presence of 0.2 M NaCl and 2.5 volumes of ethanol and then quantitated by comparison with a molecular weight standard using agarose gel electrophoresis.

5 The aptazyme construct was transcribed in a 10  $\mu$ l high yield transcription reaction (AmpliScribe from Epicentre, Madison, WI. The reaction contained 500 ng PCR product, 3.3 pmoles of P<sup>32</sup> [ $\alpha$ -32 P]UTP (3000 Camel), 1X AmpliScribe transcription buffer, 10 mM DTT, 7.5 mM each NTP, and 1  $\mu$ l AmpliScribe T7 polymerase mix. The transcription reaction was incubated at 37° C for 2 hours. One unit of RNase free-DNase was added and the reaction returned to 37° C for 30 minutes. The transcription was then purified on a 6% denaturing polyacrylamide gel to separate the full length RNA from incomplete transcripts and spliced products, eluted and quantitated spectrophotometrically.

10  
15

Sub B 14  
Cont'd

#### In vitro Assay

20 The RNA (4 pmoles/12  $\mu$ l H<sub>2</sub>O) was heated to 94° C for 1 minute then cooled to 37° C over 2 minutes in a thermocycler. The RNA was divided into 2 splicing reactions (9  $\mu$ l each) containing 100 mM Tris-HCl (pH 7.45), 500 mM KCl and 15 mM MgCl<sub>2</sub>, plus or minus theophylline (2 mM). The reactions were immediately placed on ice for 30 minutes. GTP (1 mM) was

added to the reactions (final volume of 10  $\mu$ l) and the reactions were incubated at 37° C for 2 hours.

5  
Sub B15

The reactions were terminated by the addition of stop dye (10  $\mu$ l) (95% formamide, 20 mM EDTA, 0.5% xylene cyanol, and 0.5% bromophenol blue). The reactions were heated to 70° C for 3 minutes and 10  $\mu$ l was electrophoresed on a 6% denaturing polyacrylamide gel. The gel was dried, exposed to a phosphor screen and analyzed using a Molecular Dynamics Phosphorimager (Sunnyvale, CA).

10  
Activation was determined from the amount of circular intron in each reaction. Circularized introns migrate slower than linear RNA and can be seen as the bands above the dark bands of linear RNA in the +Theo lanes of the gels of Figs. 2a and. 2b.

15  
In vivo Screening of Group I Aptazymes

20  
The aptazyme constructs as well as the wild type and a negative control were ligated into a vector that contains the T4 td intron with Eco RI and Spe I flanking the P6 region, transformed and minipreped. The plasmids were then transformed into C600:Thy A Kan<sup>R</sup> cells (cells lacking thymidine synthetase).

Individual colonies were picked and grown in rich media overnight. Theophylline (1  $\mu$ l: 6.6 mM) or H<sub>2</sub>O (1  $\mu$ l) was

added to 2  $\mu$ l of the overnight growth and was spotted on either minimal media plates, or minimal media plates with thymine.

5

## Example 2: GpIP6Thpool

### *In vitro* Selection of Group I Aptazymes

Example 2 illustrates how to generate a population of aptazymes so that there is variation in the nucleotide sequence of the aptamers. This example also illustrates how to select for phenotypes that are responsive to an effector molecule from among that population of aptazymes.

#### Construction of Pool

The construction of the pool was similar to the construction of the individual engineered aptazyme constructs. Oligos GpIWt3.129 and GpIThP6pool: 5'-GCC TGA GTA TAA GGT GAC TTA TAC TAG TAA TCT ATC TAA ACG GGG AAC CTC TCT AGT AGA CAA TCC CGT GCT AAA TN(1-4)A TAC CAG CAT CGT CTT GAT GCC CTT GGC AGN(1-4) TAA ATG CCT AAC GAC TAT CCC TT-3' (SEQ ID NO:5) were extended in the same manner as above. The extension reaction was diluted and used as template for a PCR reaction. The PCR reaction was similar to the reaction described with the following exceptions: the volume was doubled and GpIWt4.89 was replaced with GpIMutG.101: 5'-CTT AGC TAC AAT ATG AAC TAA CGT

AGC ATA TGA CGC AAT ATT AAA CGG TAG TAT TAT GTT CAG ATA AGG  
TCG TTA ATC TTA CCC CGG AAT TCT ATC CAG CT-3' (SEQ ID NO:6) in  
which there is an G to A mutation at the terminal residue of  
the intron. The pool had a diversity of  $1.16 \times 10^5$  molecules.  
5 RNA was made as described above.

In vitro Negative Selection

10 The RNA (10 pmoles/70  $\mu$ l H<sub>2</sub>O) was heated to 94° C for 1  
minute then cooled to 37° C over 2 minutes in a thermocycler.  
The splicing reaction (90  $\mu$ l) contained 100 mM Tris-HCl (pH  
7.45), 500 mM KCl and 15 mM MgCl<sub>2</sub>. The reaction was  
immediately placed on ice for 30 minutes. GTP (1 mM) was  
added to the reaction (final volume of 100  $\mu$ l) and the  
reaction was incubated at 37° C for 20 hours. The reaction  
was terminated by the addition 20 mM EDTA and precipitated in  
15 the presence of 0.2 M NaCl and 2.5 volumes of ethanol. The  
reaction was resuspended in 10  $\mu$ l H<sub>2</sub>O and 10  $\mu$ l stop dye and  
heated to 70° C for 3 minutes and was electrophoresed on a 6%  
denaturing polyacrylamide gel with Century™ Marker ladder  
(Ambion, Austin, TX). The gel was exposed to a phosphor  
20 screen and analyzed. The unreacted RNA was isolated from the  
gel, precipitated and resuspended in 10  $\mu$ l of H<sub>2</sub>O.

### Positive Selection

5 The RNA (5  $\mu$ l of negative selection) was heated to 94° C for 1 minute then cooled to 37° C over 2 minutes in a thermocycler. The positive splicing reaction (45  $\mu$ l) contained 100 mM Tris-HCl (pH 7.45), 500 mM KCl, 15 mM MgCl<sub>2</sub> and 1mM theophylline. The reaction was immediately placed on ice for 30 minutes. GTP (1 mM) was added to the reaction (final volume of 50  $\mu$ l) and the reaction was incubated at 37° C for 1 hour. The reaction was terminated by the addition of stop dye, heated to 70° C for 3 minutes and was electrophoresed on a 6% denaturing polyacrylamide gel with Century™ Marker ladder. The gel was exposed to a phosphor screen and analyzed. The band corresponding to the linear intron was isolated from the gel and precipitated and resuspended in 20  $\mu$ l H<sub>2</sub>O.

### Amplification and Transcription

10  
15  
20 The RNA was reverse transcribed in a reaction containing 250 mM Tris-HCl (pH 8.3), 375 mM KCl, 15 mM, MgCl<sub>2</sub>, 0.1 M DTT, 0.4 mM of each dNTP 2  $\mu$ M GpIMutG.101 and 400 units of SuperScript II reverse transcriptase (Gibco BRL, Rockville, MD). The cDNA was then PCR amplified, transcribed and gel purified as described above.

Figure 3 depicts an in vivo assay system for Group I introns of the present invention. The td intron normally sits within the td gene for thymidylate synthase (TS) in phage T4. A *ThyA E. coli* host that lacks cellular TS is unable to grow in the absence of exogenous thymine or thymidine (-Thy). The cloned td gene can complement the *ThyA* cells and grow on -Thy media. Conversely, cells that lack TS have a selective advantage on media containing thymidine and trimethoprim. Therefore, cells harboring theophylline-responsive Group I aptazymes grow better in the presence of theophylline and the absence of thymidine. In contrast, the same cells grow better in the absence of theophylline and the presence of thymidine and trimethoprim.

This strategy provides both a positive in vivo screen and selection for theophylline-dependent activation and a negative in vivo screen and selection for theophylline-absent repression. The assay system of Fig. 3 was used in Example 1, above, for the in vivo screening of Group I aptazymes in a specific embodiment of the present invention.

Figure 4a depicts the critical residues of the P6 region of the Group I ribozyme joined to the anti-theophylline aptamer by a short randomized region to generate a pool of aptazymes of the present invention. The residues shown in

bold in Fig. 4a are the P6 critical residues, and the faded residues shown in Fig. 4a are the anti-theophylline aptamer. The randomized regions are designated in Fig. 4a as "N1-4". Approximately 40 random sequence residues are introduced into the N1-4 region of the construct to synthesize a pool of aptazymes, referred to herein as a communication module pool.

Figure 4b shows a selection protocol for the Group I P6 Aptazyme Pool of Fig. 4a. Positive and negative selections are made in vitro to select Group I aptazymes that are dependent on theophylline. The selections are described above in Example 2 for a specific embodiment of the present invention. In vivo screens and selections are used to select Group I aptazymes that exhibit strong theophylline-dependence. The selected aptazymes are mixed at various ratios with mutant Group I ribozymes that splice at a low but continuous level to determine the level at which aptazymes can be selected against background. A communication module pool can be transformed with the selected aptazymes to determine whether the same modules that function in vitro are also functional in vivo. Finally, the best theophylline-dependent Group I aptazymes that have been derived by any of the methods described herein may undergo further selection by partially randomizing their sequences and selecting for improved in vivo performance.



Strategies similar to those depicted in Figs 4a and 4b may be used to develop aptazymes and aptamers dependent on any desired effector molecule. See generally G.A. Soukup, et al., Engineering precision RNA molecular switches. *Proc. Natl. Acad. Sci. U.S.A.* **96**, 3584-3589 (1999) and M. Koizumi, et al., Allosteric selection of ribozymes that respond to the second messengers cGMP and cAMP. *Nature Struct. Biol.* **6**, 1062-1071 (1999). Positive and negative in vitro selection such as depicted in Fig. 4b are described above in Example 2 for a specific embodiment of the present invention.

From  $10^6$  to  $10^{10}$  variants can be efficiently transformed as described herein, sufficient to encompass most variants in the populations discussed so far. This efficiency of transformation, however, is likely to be insufficient to encompass a significant fraction of a completely random pool. Nonetheless, sequences have been selected from completely random expressed pools that can protect bacteria from bacteriophage infection.

The optimization strategies described herein yield Group I aptazymes that are highly dependent on small molecule effectors. Since the rules that govern Group I intron splicing in different gene contexts are well known to those skilled in the art, the skilled artisan can remove Group I

aptazymes from the td gene and modularly insert them into other genes. Should the efficiency or effector-dependence of intron splicing be compromised in the new gene, the intron can be reaccommodated to its new genetic environment by fusing td or another selectable marker to the interrupted gene of interest and selecting for an effector-dependent phenotype.

To the extent that Group I aptazymes are self-sufficient, they should also function in eukaryotic cells, including human cells. However, to the extent that the architecture of eukaryotic cells is significantly different from the cytoplasm of bacteria, the efficiency or effector-dependence of intron splicing may suffer on aptazyme transfer between different species of organisms. Selecting for effector-dependence, but now in eukaryotic cells, may be necessary to obtain satisfactory efficiency and effector-dependence. Selection in eukaryotic systems may be performed by fusing the gene of interest to a reporter gene such as GFP or luciferase. Variants of the aptazyme that promote effector-dependent protein production may then be selected using a FACS.  $10^6$  to  $10^8$  variants may be screened by this procedure, a range comparable to the bacterial system previously described.

Figure 5 is a diagrammatic representation of one embodiment of the exogenous or endogenous activation of Group

5  
Sub B19  
Contd

I intron splicing is depicted. A gene of interest 10 is fused to a reporter gene 12 such as luciferase or beta-galactosidase, which also contains the group I intron (td) 14. Splicing-out of the Group I intron is induced by an endogenous effector molecule 16, which may be a protein, e.g., Cyt18. Alternatively, splicing-out of the Group I intron may be induced by an exogenous effector molecule 18. Activation of the aptazyme and auto-excision of the intron results in expression of the reporter gene encoded protein 20 that is detect by, e.g., fluorescence 22 or any other desired detectable reaction. The use of a reporter gene 12 of this embodiment may be suitable for use in eukaryotic systems.

10  
15  
20

Figure 6 is a diagram of another embodiment of the present invention. Libraries of candidate exogenous activators 30 can be generated from a randomized aptazyme pool indicated by random loop  $E_{1-n}$ . As in the embodiment of Fig. 5, a reporter gene 12 is expressed in cells where the exogenous activator 30 activates the aptazyme to release the intron, which may contain a random loop 32, from the gene. In this embodiment, the reaction occurs within cells which are then sorted 34 based on a chromogenic reaction or emission 22, or may even be isolated by, e.g, statistical cell separation cloning. As will be known to those of skill in the art of

enzymatic oligonucleotides any number of current and future effector molecule libraries may be used with the present invention.

Figure 7 depicts an alternative embodiment for screening libraries of exogenous activators. In the embodiment of the present invention of Fig. 7, Group I introns with length polymorphisms are induced into the construct by trans-splicing with an independent oligonucleotide. Libraries of candidate exogenous activators 30 can be generated from a randomized aptazyme pool indicated by random loop  $E_{1-n}$ . As in the embodiment of Fig. 5, a reporter gene 12 is expressed in cells where the exogenous activator 30 activates the aptazyme to release the intron, which may contain a random loop 32, from the gene. In this embodiment, the reaction occurs within the intron 14 and an independent oligonucleotide 36 by a trans-splicing reaction and extraction step 38. Extracted trans-spliced intron reporter gene constructs are then amplified by, e.g., polymerase chain reaction in step 40, followed by transformation of cells with the transplined construct at step 42. Transformation of the transplined construct may be performed by those of skill in the art with either a negative or positive selection scheme for identification of the trans-spliced gene.

Figure 8 shows yet another alternative embodiment for screening libraries of exogenous activators 30 with the present invention. In the embodiment of Fig. 8, a pool of randomized loops 32 interact with an effector 16 (or a second exogenous effector) to form a complex 44. The complex 44 is exposed to the gene 10 containing both the Group I intron 14 and a reporter gene 12. Cell sorting 34 reveals the cells that express the reporter gene 20 to indicate successful activation of the aptazyme by the effector.

Figure 9 is a diagram of an embodiment of the present invention for screening for endogenous suppressors. In this embodiment, an endogenous effector 46, in this illustration shown as a protein suppressor from endogenous or transformed origin, activates self-splicing of the Group I intron 14 (depicted with a random loop 32). Cell sorting 34 is used to reveal the expression of the reporter gene. To protect against spontaneous auto-excision of the intron, the gene may be transferred into a different background system (at step 48) such as yeast or *E. coli*, for example.

Figure 10 depicts an alternative to the embodiment of Fig. 9 to screen for endogenous activators of the present invention. In Fig. 10, the activator that is being screened for may include, *inter alia*, a phosphorylated protein, a

product of ubiquitination, or a protein-protein complex. For example, a protein suppressor 46, may phosphorylate 50 an effector molecule 16 (e.g., Cyt18). The phosphorylated effector molecule 52 activates intron 14 self-splicing with concomitant expression of the reporter gene 12, e.g., green fluorescent protein (GFP).

Figure 11 shows yet another embodiment of the present invention to screen for compounds that perturb cellular metabolism. In this embodiment, a communication module pool 54, undergoes selection for a phenotype responsive to a protein suppressor 46 effector molecule. The protein effector 46 may be a phosphoprotein, an induced protein, or a protein complex, for example. The source of the effector may be endogenous, exogenous or may even be the product of a transformation construct used to transform a cell. Activation by the effector results in expression of the reporter gene 12, but may inactivate, suppress, or "knocks-out" the gene 10 of interest. The functioning of the gene of interest may thereby be perturbed, providing information about the function and/or regulation of the gene or gene product.

Figure 12 shows a further embodiment of the present invention that provides a non-invasive readout of metabolic states. An aptazyme construct of the present invention may be

introduced into a gene of interest 56 that forms part of the host (whether chromosomal or extrachromosomal). A protein suppressor 46 from either an endogenous or exogenous source is used to screen for the product of a cell transformation that may activate self-splicing of the aptazyme, leading to expression of the reporter gene 10. Whether or not the gene of interest is expressed upon release of the aptazyme intron from the gene provides information about the metabolic state of the gene of interest. The embodiment of the present invention of Fig. 12 thus, provides a non-invasive means to determine the metabolic state of an organism with regard to a gene of interest.

Figure 13 depicts a further embodiment of the present invention wherein endogenous suppressors provide a non-invasive readout of multiple metabolic states. Multiple protein suppressors 46 (endogenous or transformed) are exposed to a pool of Group I introns 14 of the present invention. The pool includes introns with length polymorphisms that are depicted in Fig. 13 by a discontinuity or break in the line representing the Group I intron 14 residing in a gene of interest 10. Activation of the aptazyme leads to trans-splicing among the various polymorphisms 58. The products of trans-splicing may be extracted and amplified in step 60.

Separation of the trans-splicing products 58 by gel electrophoresis 62 provides a read out of the protein function or the metabolic pathway. The readout may even be digitized for analysis.

5        Figure 14 depicts schematically an exemplary worksurface for yet another embodiment of the present invention: automated selection. See, J.C. Cox, et al., *Automated RNA Selection* Biotechnol. Prog., 14, 845 850, 1998.

10        Base protocol. Automated selection involves several, sequential automated steps. Several modules are placed on the robotic worksurface, including a magnetic bead separator, and enzyme cooler, and a thermal cycler. After manually preparing reagents and preloading those reagents (including random pool RNA, buffers, enzymes, streptavidin magnetic beads, and biotinylated target) and tips onto the robot, a program is run. The selection process, automated by the robot, goes as follows: RNA pool is incubated in the presence, of biotinylated target conjugated to streptavidin magnetic beads. After incubation, the magnets on the magnetic bead separator are raised, and the beads (now bound by pool RNA - the selected aptamers) are pulled out of solution. Thus, the beads can be washed, leaving only RNA bound to targets attached to beads.

15

20



These RNA molecules are reverse transcribed, reamplified via PCR, and the PCR DNA is in vitro transcribed into RNA to be used in iterative rounds of selection.

The Bioworks method for in vitro selection. This  
5 scripted programming method contains all movements necessary in order to facilitate automated selection. This includes all physical movements to be coordinated, and also communication statements. For instance, five rounds of automated selection against a single target requires over 5,000 separate movements  
10 in x, y, z, t coordinate space. Additionally, the method also holds all relevant measurements, offsets, and integrated equipment data necessary to prevent physical collisions and permit concerted communication between devices.

"Beads on filter" selections. While the vast majority of  
15 manual selections have been performed on nitrocellulose-based filters, a small few have also been performed on solid surfaces, such as beads. We have developed a novel selection scheme whereby selection is performed on magnetic beads that are placed on nitrocellulose filters, and washed as the bead  
20 is the selection target itself. This method allows for much greater specificity of selection, thereby promoting 'winning' molecules to amplify in greater number, and thus reduce the overall amount of rounds necessary to complete the selection

procedure. Manual selection does not involve a combination of surfaces to enhance selection.

Cross-contamination avoidance. The introduction of contaminating species of nucleic acid strands in a manual selection can be commonplace. This is especially true if selection is done against multiple targets in parallel, and also when a researcher reuses the same pool for different selection tools. Contaminating species have been shown in the past to interfere with a manual selection such that it could not be completed. Automated *in vitro* selection takes steps to minimizing and/or eliminate the possibility of cross-contamination between pools and targets. Movement of the mechanical pod along the worksurface is unidirectional when carrying potentially contaminating material. This movement away from 'clean' things and only towards items that have already been exposed to replicons greatly diminishes the possibility of cross-contaminating reactions. The only circumstance in which the pod reverses its direction is to acquire a new, clean pipette tip. Additionally, we have also sealed our reagent trays with aluminum foil for a physical barrier between the environment and unexposed reagents. See Figure 14, a layout of the robotic worksurface which reduces cross-contamination.

Figure 15a depicts the LI ligase used for pool design in, e.g., the Cyt18 aptazyme selection, as an example of a protein-activated aptazyme. Stems A, B, and C are indicated. The shaded region contains the catalytic core and ligation junction. Primer binding sites are shown in lower case, an oligonucleotide effector required for activity is shown in italics, and the ligation substrate is bolded. The 'tag' on the ligation substrate can be varied, but was biotin in the exemplary selection described herein. The LI pool contains 50 random sequence positions and overlaps with a portion of the ribozyme core. In Figure 15b, Stem B was reduced in size and terminated with a stable GNRA tetraloop, but stem A was unchanged.

Figure 15c schematically shows the following selection scheme: the RNA pool was incubated with a biotinylated tag and reactive variants were removed from the population. The remaining species were again incubated with a biotinylated tag in the presence of the target (Cyt18). Reactive variants were removed from the population and preferentially amplified by reverse transcription, PCR, and in vitro transcription.

Figures 16a through 16c schematically depict one method to anchor aptazymes. A substrate 70, e.g, glass, silicon, gallium arsenide, silicon on insulator (SOI) structures,

epitaxial formations, germanium, germanium silicon, polysilicon, amorphous silicon, and/or like substrate, semi-conductive or conductive is depicted. The substrate 70 may already have been processed to provide electrical means of detection on the surface of the wells 72. One example of such a detector is a charge coupled capacitor.

In Fig. 16a, different ribozyme ligases 74 are shown immobilized on beads 73 in wells 72, and mixtures of analytes 76 and tagged substrates 78 may be added to each well. Next, in Fig. 16b, cognate effectors are present (same analyte and allosteric site) in the well 72 and the aptazymes 74 will covalently attach the reporter tags 80 (e.g., fluorescent tags) to themselves. Where aptazymes 74 have not been activated by effectors, the tagged substrates 78 are washed out of the well. In Fig. 16c, after reaction and washing, the presence and amounts of co-immobilized reporter tags 80 are indicative of amounts of ligands that were present during the reaction. See K. A. Marshall, et al., Training Ribozymes to Switch, *Nature Struct. Biol.* 6 (11) 992-994, 1999.

In the embodiment of Figure 16, the reporter tag 80 may be an enzyme, a magnetic particle, or any number of detectable particles. Additionally, the ribozymes could be immobilized

on beads 83, but they could also be directly attached to a solid support or substrate 70 via covalent bonds.

One advantage of this embodiment is that covalent immobilization of reporters allows stringent wash steps to be employed. This can be distinguished from non-covalent immobilization assays such as ELISA™ assays where stringent washing may destroy the signal. An additional advantage is that ribozyme ligases have the unique property of being able to transduce effectors into templates that can be amplified, affording an additional boost the in signal prior to detection.

Although nucleic acids are generally less robust than antibodies, modified nucleotides may be introduced into the aptazymes that substantially stabilize them from degradation in environments such as sera or urine. Antibodies generally have higher affinities for analytes than do aptamers. However, the analytical methods of the present invention do not rely on binding per se, but only on transient interactions. The present invention requires mere recognition rather than actual binding, thus providing a potential advantage of aptazymes over antibodies. That is, even low affinities are sufficient for activation and subsequent detection, especially if individual immobilized aptazymes are

examined (i.e., by ligand-dependent immobilization of a quantum dot).

All publications mentioned in the above specification are hereby incorporated by reference. Modifications and variations of the described compositions and methods of the invention will be apparent to those skilled in the art without departing from the scope and spirit of the invention'. Although the invention has been described in connection with specific embodiments, it should be understood that the invention as claimed should not be unduly limited to such specific embodiments. Indeed, various modifications of the described compositions and modes of carrying out the invention which are obvious to those skilled in molecular biology or related arts are intended to be within the scope of the following claims.